

Ethanol potently and competitively inhibits binding of the alcohol antagonist Ro15-4513 to $\alpha_4/\beta_3\delta$ GABA_A receptors

H. Jacob Hancher^{*†}, Panida Chutrinopkun^{*‡}, Pratap Meera[§], Porntip Supavilai[†], Werner Sieghart[¶], Martin Wallner^{*¶}, and Richard W. Olsen^{*¶}

Departments of ^{*}Molecular and Medical Pharmacology and [§]Neurobiology, University of California, Los Angeles, CA 90095; [†]Department of Pharmacology, Faculty of Science, Mahidol University, Bangkok 10 400, Thailand; and [¶]Center for Brain Research, Division of Biochemistry and Molecular Biology and Section of Biochemical Psychiatry, Medical University of Vienna, A-1090 Vienna, Austria

Edited by Floyd E. Bloom, The Scripps Research Institute, La Jolla, CA, and approved February 10, 2006 (received for review November 16, 2005)

Although GABA_A receptors have long been implicated in mediating ethanol (EtOH) actions, receptors containing the “nonsynaptic” δ subunit only recently have been shown to be uniquely sensitive to EtOH. Here, we show that δ subunit-containing receptors bind the imidazo-benzodiazepines (BZs) flumazenil and Ro15-4513 with high affinity ($K_d < 10$ nM), contrary to the widely held belief that these receptors are insensitive to BZs. In immunopurified native cerebellar and recombinant δ subunit-containing receptors, binding of the alcohol antagonist [³H]Ro15-4513 is inhibited by low concentrations of EtOH ($K_i \approx 8$ mM). Also, Ro15-4513 binding is inhibited by BZ-site ligands that have been shown to reverse the behavioral alcohol antagonism of Ro15-4513 (i.e., flumazenil, β -carbolinecarboxylate ethyl ester (β -CCE), and *N*-methyl- β -carboline-3-carboxamide (FG7142), but not including any classical BZ agonists like diazepam). Experiments that were designed to distinguish between a competitive and allosteric mechanism suggest that EtOH and Ro15-4513 occupy a mutually exclusive binding site. The fact that only Ro15-4513, but not flumazenil, can inhibit the EtOH effect, and that Ro15-4513 differs from flumazenil by only a single group in the molecule (an azido group at the C7 position of the BZ ring) suggest that this azido group in Ro15-4513 might be the area that overlaps with the alcohol-binding site. Our findings, combined with previous observations that Ro15-4513 is a behavioral alcohol antagonist, suggest that many of the behavioral effects of EtOH at relevant physiological concentrations are mediated by EtOH/Ro15-4513-sensitive GABA_A receptors.

alcohol receptor | flumazenil | β -carbolines | extrasynaptic GABA_A receptors

Although many proteins show changes in their function at very high alcohol concentrations (>50 mM), the molecular basis for behavioral alcohol effects at low to moderately intoxicating doses experienced during social alcohol consumption remains elusive (1). GABA_A receptors (GABA_ARs) and the inhibitory GABAergic system have long been suspected to be targets for acute alcohol effects (2–4). For example, the GABA_AR agonist muscimol potentiates the sedative actions of alcohol, whereas the opposite effect, a reduction of ethanol (EtOH)-produced sedation, is detected with the GABA_AR blockers picrotoxin and bicuculline (5). Although most GABA_AR subunit combinations can be activated by high (anesthetic) alcohol concentrations (6), only very specific GABA_AR subunit combinations (containing the δ as well as the β_3 subunit) exhibit dose dependencies that mirror blood alcohol levels that are associated with mild to moderate intoxication in humans (7, 8) (≈ 3 –30 mM, because the legal drinking limit is 17 mM or 0.08%). GABA_ARs containing the δ subunit are located either outside (9) or in the perimeter of (10) synapses but not in the subsynaptic membrane, and they give rise to a persistent (tonic) GABA current (11) that is enhanced by low alcohol concentrations (12–14). However, there is controversy because Carta *et al.*

(14) concluded that the EtOH-induced increase in tonic current is due to increased GABA release rather than a postsynaptic effect. Also, for reasons on which we can only speculate, a recent article by Borghese *et al.* (15) states that effects of low dose EtOH on δ subunit-containing receptors could not be observed, in particular with human $\alpha_4\beta_3\delta$ GABA_AR clones expressed in oocytes and cell lines.

EtOH pharmacology shares many characteristics with allosteric activators of GABA_ARs (loosely referred to as GABA_AR agonists), such as benzodiazepines (BZs) (5). Additional evidence for a link between EtOH and BZ actions on GABA_AR comes from the surprising finding that a mutation in the α_6 subunit (α_6 -R100Q), which was identified in alcohol-nontolerant and -nonpreferring rats in refs. 16 and 17, leads to receptors with increased alcohol sensitivity in recombinant expression (in cerebellar granule cell neurons) and to alcohol-hypersensitive animals in behavioral studies (12). Histidine residues at positions that are homologous to the α_6 -R100 residue (which affects EtOH sensitivity in $\alpha_6\beta_3\delta$ GABA_ARs) are critical for high-affinity binding of classical BZ agonists at the interface between α and γ_2 subunits.

It has been thought that the “extrasynaptic” GABA_AR δ subunit, which presumably takes the position of the γ_2 subunit in functional pentameric GABA_ARs, renders receptors insensitive to BZs (18). Also, most δ subunits are found to be associated with α_4 and α_6 subunits that differ from other GABA_AR α subunits at a critical position (a histidine in $\alpha_{1,2,3,5}$ replaced by an arginine in $\alpha_{4,6}$) that makes $\alpha_4/\beta_3\gamma_2$ receptors insensitive to classical BZ agonists such as diazepam (DZ) and flunitrazepam (19, 20). However, Arg-100 (WT) in α_4 and α_6 receptors still allows high-affinity binding of the imidazo-BZs Ro15-4513 and flumazenil (in $\alpha_6\beta_3\gamma_2$ GABA_ARs). In functional receptor assays, Ro15-4513 is a weak partial inverse agonist (i.e., it leads to a slight reduction in GABA_AR activity) on the most abundant GABA_AR subtypes in the brain (20). Ro15-4513 is a partial agonist (i.e., it enhances GABA action, but less than DZ even at saturating concentrations) on the α_4 and α_6 receptors (with β and γ_2 subunits), whereas flumazenil is essentially silent in functional assays. Interestingly, Ro15-4513, but not other inverse agonists

Conflict of interest statement: M.W., R.W.O., and H.J.H. have filed a U.S. Provisional Patent Application, Serial No. 60/693,844.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: GABA_AR, GABA_A receptor; HEK, human embryonic kidney; BZ, benzodiazepine; EtOH, ethanol; IP, immunoprecipitated/immunoprecipitation; β -CCE, β -carboline-3-carboxyethyl ester; DMCM, methyl-6,7-dimethoxy-4-ethyl- β -carboline-3-carboxylate; DZ, diazepam; DZ-15, DZ-insensitive; FG7142, *N*-methyl- β -carboline-3-carboxamide.

[†]H.J.H. and P.C. contributed equally to this work.

[¶]To whom correspondence may be addressed at: Department of Molecular and Medical Pharmacology, University of California, Room 23-120 CHS, Charles Young Drive South, Los Angeles, CA 90095-1735. E-mail: mwallner@mednet.ucla.edu or rolsen@mednet.ucla.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 3. Test for competitive antagonism between EtOH and Ro15-4513. Ro15-4513 saturation binding was performed in the presence of 10, 30, and 50 mM EtOH. Individual curves were fitted with the Hill equation (shown as black curves). Colored curves are derived from a simultaneous least-square fit of the entire data set by using the following equation: $[^3\text{H}]\text{Ro15-4513} = (f/(1-f)^{1/h}) \times K_d \times (1 + [\text{EtOH}]/K_{d\text{EtOH}})$. $[^3\text{H}]\text{Ro15-4513}$ is the concentration of $[^3\text{H}]\text{Ro15-4513}$ to reach a fractional occupancy (f), h is the Hill coefficient, K_d is the dissociation constant for $[^3\text{H}]\text{Ro15-4513}$, and the term $(1 + [\text{EtOH}]/K_{d\text{EtOH}})$ is derived from the Schild–Gaddum equation and describes the decreased receptor occupancy by Ro15-4513 in the presence of the proposed competitive antagonist EtOH, where $[\text{EtOH}]$ is the EtOH concentration, and $K_{d\text{EtOH}}$ is the dissociation constant for EtOH. The best fit of curves was obtained with a Hill coefficient of 1.1, a $K_{d\text{EtOH}}$ of 8.0 mM, and a K_d for Ro15-4513 of 7.5 nM. (Inset) Dissociation of $[^3\text{H}]\text{Ro15-4513}$ from $\alpha_4\beta_3\delta$ receptors equilibrated with 5 nM $[^3\text{H}]\text{Ro15-4513}$. The unbinding rate of $[^3\text{H}]\text{Ro15-4513}$ was measured by adding excess (1 μM) cold Ro15-4513 followed by rapid filtration after ≈ 0.5 , 1, 2, 5, and 10 min, and counting the amount of bound hot ligand. To test the effect of EtOH on the dissociation rate, 200 mM EtOH was added instead of cold Ro15-4513 to prevent rebinding of the radioligand after dissociation. This experiment is representative for a total of three experiments that were performed.

ies (29), $\approx 20\%$ of the $[^3\text{H}]\text{Ro15-4513}$ binding to cerebellar membranes was not blocked by 10 μM DZ (Fig. 2b Inset). Of this DZ-insensitive $[^3\text{H}]\text{Ro15-4513}$ binding, $\approx 30\%$ was dose-dependently inhibited by EtOH ($\text{IC}_{50} \approx 7$ mM), with a maximum inhibition at 100 mM EtOH (Fig. 2b). Given that DZ-IS binding of $[^3\text{H}]\text{Ro15-4513}$ in the cerebellum is $\approx 20\%$ of total binding, EtOH-displaceable Ro15-4513 binding is $\approx 6\%$ of total cerebellar Ro15-4513 binding. This percentage is consistent with the δ Ab-precipitated fraction, enriched in $\alpha_6\beta\delta$ receptors and in the same range as the fraction of $\alpha_6\beta_x\delta$ receptors ($\approx 11\%$) determined by biochemical methods in rat cerebellum (30). EtOH did not inhibit the $[^3\text{H}]\text{Ro15-4513}$ binding to classical DZ-sensitive sites under the same conditions (data not shown).

Ro15-4513 Is a Competitive Alcohol Antagonist. The complete and dose-dependent displacement of $[^3\text{H}]\text{Ro15-4513}$ by EtOH on recombinant $\alpha_4\beta_3\delta$ receptors (Fig. 2a) suggests the possibility of a competitive antagonism between EtOH and Ro15-4513. To evaluate and distinguish competitive (direct) from allosteric (indirect) interaction further, we performed $[^3\text{H}]\text{Ro15-4513}$ saturation-binding experiments in the continuous presence of nonsaturating EtOH concentrations. Receptor occupancy of ligands (in our case, $[^3\text{H}]\text{Ro15-4513}$) decreases in a predictable way in the presence of a presumed competitive antagonist (EtOH). The inclusion of 10, 30, or 50 mM EtOH in $[^3\text{H}]\text{Ro15-4513}$ saturation-binding experiments led to a dose-dependent parallel shift of the $[^3\text{H}]\text{Ro15-4513}$ binding curve to the right (Fig. 3). The simultaneous least-square fit using a combined Hill/Schild–Gaddum equation (with $K_d = 7.5$ nM for Ro15-4513) resulted in a Hill coefficient of 1.1 and a $K_d = 8.1$ mM for




Fig. 4. Pharmacological characterization of the $[^3\text{H}]\text{Ro15-4513}$ -binding site. Displacement of 10 nM $[^3\text{H}]\text{Ro15-4513}$ by the BZ antagonist flumazenil (Ro15-1788), the Ro15-4513 congeners RY080 and RY024, and the BZ site ligands and β -CCE on recombinant $\alpha_4\beta_3\delta$ receptors expressed in HEK cells. Very similar results were obtained displacing $[^3\text{H}]\text{Ro15-4513}$ with flumazenil, FG7142, and β -CCE from native immunopurified δ subunit-containing cow cerebellar GABA_ARs (data not shown).

EtOH (colored curves in Fig. 3). That the prediction for competitive antagonism overlaps with our experimental data provides additional evidence that alcohol and Ro15-4513 are likely to occupy overlapping binding sites.

The decrease in receptor occupancy in the presence of competitive antagonists, as described by the Schild–Gaddum equation, is due to a reduction in the apparent association rate of ligands (binding sites occupied by the competitive ligand are not available for binding), without changes in the dissociation rate (i.e., the residence time of ligands in their binding sites). Therefore, we decided to determine the dissociation rate of $[^3\text{H}]\text{Ro15-4513}$ from recombinant $\alpha_4\beta_3\delta$ receptors by measuring unbinding after the addition of excess cold (1 μM) Ro15-4513 or a high concentration of EtOH (200 mM) to a receptor preparation equilibrated with 5 nM $[^3\text{H}]\text{Ro15-4513}$. Excess (1 μM) Ro15-4513 as well as 200 mM EtOH led to essentially identical $[^3\text{H}]\text{Ro15-4513}$ ligand dissociation rates, showing that EtOH does not decrease the residence time of $[^3\text{H}]\text{Ro15-4513}$ in its receptor (Fig. 3 Inset). This result suggests that EtOH, even at 200 mM, does not allosterically alter the binding pocket.

Ro15-4513 Binding to $\alpha_4\beta_3\delta$ GABA_ARs Is Inhibited by Ligands That Prevent the Behavioral Alcohol Antagonism of Ro15-4513. To further characterize the $[^3\text{H}]\text{Ro15-4513}$ binding site on the δ subunit receptors, we performed competition binding experiments on recombinant ($\alpha_4\beta_3\delta$) and native (cow cerebellum) immunopurified δ subunit-containing receptors with BZ site ligands that have been shown to reverse the behavioral alcohol antagonism of Ro15-4513 [flumazenil, β -CCE, and *N*-methyl- β -carboline-3-carboxamide (FG7142)] (22, 31) and the imidazo-BZs (Ro15-4513/flumazenil structural analogs) RY024 and RY080. RY024 has been described as a behavioral alcohol antagonist (32, 33). We show that like “cold” Ro15-4513, flumazenil, RY024, and RY080, and the β -carbolines β -CCE and FG7142, displaced $[^3\text{H}]\text{Ro15-4513}$ from its binding site on $\alpha_4\beta_3\delta$ receptors (Fig. 4). Therefore, ligands that block EtOH enhancement of $\alpha_4/\alpha_6\beta_3\delta$ GABA_AR currents, such as Ro15-4513, or that reverse the alcohol-antagonist activity of Ro15-4513 (flumazenil and β -CCE) (21, 29, 45), are also able to displace $[^3\text{H}]\text{Ro15-4513}$ binding. However, most other BZ site ligands [including all of the tested classical BZ agonists (DZ, flurazepam, flunitrazepam, and midazolam), the β -carboline DMCM, and the BZ site ligands zolpidem and zopiclone] that are known to bind with high affinity to the classical BZ sites in γ_2 subunit-containing recep-

[³H]flumazenil binding is not displaced by EtOH (≤ 300 mM) suggest that the C7 azido group is critical for alcohol antagonism. Therefore, we propose a model in which the azido group (which is approximately the same size as alcohol) occupies the EtOH-binding pocket in the Ro15-4513-bound receptor and thus prevents EtOH binding to these types of GABA_ARs. Consistent with this view, there have been recent reports of *in vivo* alcohol-antagonist actions of the Ro15-4513/flumazenil derivative RY024; RY024 and the related molecule RY080 have an acetylene group (instead of an azido group) at the C7 position of the BZ ring (32). We found that these chemical analogs of Ro15-4513 (RY024 and RY080) are potent inhibitors of [³H]Ro15-4513 binding to this alcohol/BZ site. This inhibition suggests that the reported behavioral alcohol antagonism of RY024 may be due to reversal of alcohol actions on unique EtOH/BZ-sites on GABA_ARs.

Photolabeling, site-directed mutagenesis, cysteine-accessibility studies, and BZ pharmacophore models derived from those data lead to the consensus that side-chain residues at the C7 position of the BZ ring are close to the critical histidine residue (α_1 -H102, bovine numbering) that determines sensitivity of $\alpha_{1,2,3,5}$ -containing receptors to classical BZ agonists in $\alpha\beta\gamma_2$ receptors (36–40). The selective labeling, by the photoreactive C7 azido group of Ro15-4513, of tyrosine α_1 -Y210 (bovine numbering) (37), which is in close proximity to α_1 -H102 in structural homology models (37, 38, 41), is probably due to a higher photoreactivity of tyrosine residues (37).

The residue homologous to α_1 -H102 in α_6 is polymorphic in rats and can be α_6 -R100 or α_6 -Q100. We showed that the α_6 -100Q polymorphism, which has been selected for during breeding of alcohol hypersensitive (alcohol-nontolerant) rats (16), leads to a dramatically increased alcohol sensitivity of α_6 -100Q $\beta_3\delta$ GABA_AR *in vitro* and *in vivo* (12). Based on our findings described here, it is tempting to speculate that the reason for the increased alcohol sensitivity of α_6 -100Q $\beta_3\delta$ receptors (and α_6 -100Q/Q rats) is that the α_6 -100 residue is very close to, and might even directly line, the EtOH-binding site.

The data presented here strongly support the view that the alcohol-antagonist action of Ro15-4513 is largely due to specific actions on alcohol-sensitive subtypes (like $\alpha_{4/6}\beta_3\delta$) of GABA_ARs. It remains to be determined whether the partial inhibition (inverse agonist action) of Ro15-4513 on some GABA_AR subtypes contributes to this alcohol antagonism. However, the fact that other inverse agonists (e.g., FG7142 and β -CCE) are not alcohol antagonists (22, 24, 42) suggests that such a contribution from action on other GABA_AR subtypes is only minor. The inverse agonists FG7142 and β -CCE (like flumazenil) reverse the behavioral alcohol antagonism of Ro15-4513 (22, 31) and displace Ro15-4513 from the EtOH/Ro15-4513 site on $\alpha_{4/6}\beta_3\delta$ receptors. Although the relative contributions of individual GABA and non-GABA receptors to the various aspects of EtOH intoxication, as well as whether there are other unrecognized subtypes of GABA_ARs that are also sensitive to EtOH/Ro15-4513 that may contribute to Ro15-4513-sensitive behavioral EtOH effects, remain to be determined, the results presented here strongly support the view that EtOH/Ro15-4513-sensitive GABA_ARs are important mediators of alcohol actions at blood-alcohol concentrations that are experienced during low and moderate alcohol consumption. Also, the competitive nature of the EtOH–Ro15-4513 interaction on δ subunit-containing GABA_AR might allow us to determine where in the receptors the EtOH-binding site is located. The displacement of Ro15-4513 from $\alpha_{4/6}\beta_3\delta$ receptors provides a test-tube assay to screen for ligands that can bind to, or very close to, the EtOH-binding site that mediates behaviorally relevant low dose EtOH effects. This binding assay allows further structure–function analysis of the pharmacological potency and efficacy of compounds that are active at the “alcohol

receptor.” Ligands that are active at the alcohol/BZ binding site on δ subunit-containing GABA_AR receptors may yield therapeutics for problems that are related to low dose EtOH sensitivity, including anxiety, insomnia, and alcoholism.

Methods

Radiolabeled [³H]Ro15-4513 (ethyl 8-azido-5,6-dihydro-5-methyl-6-oxo-4*H*-imidazo[1,5-*a*]BZ-3-carboxylate (33.3 Ci/mmol; 1 Ci = 37 GBq) was purchased from PerkinElmer and was supplied dissolved in EtOH. To change the solvent from EtOH to DMSO, the [³H]Ro15-4513 EtOH solution was dried in a vacuum concentrator and redissolved in DMSO. Unlike EtOH, DMSO at final concentrations ($<1\%$) did not change [³H]Ro15-4513 binding to cerebellar δ -IP receptors (data not shown).

Ro15-4513, flumazenil (ethyl 8-fluoro-5,6-dihydro-5-methyl-6-oxo-4*H*-imidazo[1,5-*a*][1,4]BZ-3-carboxylate), DZ, and flunitrazepam were obtained from Hoffman–LaRoche; DMCM was a gift from Ferrosan (Copenhagen); and FG7142 and β -CCE were provided by Schering. EtOH, GABA, and bicuculline were purchased from Sigma. Compounds were dissolved in DMSO as a 10 mM stock solution and used at the indicated concentrations. Protein concentration was determined with the BCA protein assay kit (Pierce) with BSA as standard.

Membrane Preparation. Bovine cerebellum was obtained from a local supplier and stored frozen at -70°C . Tissue was thawed and homogenized by sonication in 10 volumes of homogenization buffer [50 mM Tris-HCl, pH 8.0/50 mM KCl/1 mM EDTA/0.32 M sucrose/0.5 mM DTT/0.01% bacitracin supplemented with either protease inhibitors (2 mM benzamide/0.1 mM benzethonium chloride/0.3 mM PMSF) or a protease inhibitor mixture (Complete Mini, Roche Diagnostics)] and centrifuged ($550 \times g$) for 10 min at 4°C to pellet nuclei and cells. The supernatant fraction was collected by three sonication–centrifugation cycles (in homogenization buffer without sucrose). Membrane pellets were collected at $35,000 \times g$ for 1 h at 4°C and used for ligand binding, or they were stored frozen at -20°C .

Recombinant Cell Expression. HEK 293T cells were transfected with rat cDNAs under the control of a CMV promoter ($\alpha/\beta/\delta$ or γ_2 , 1:1:2) as described in ref. 43, and cells were harvested at 60–100 h after transfection. Membranes from these HEK 293T pellets were homogenized by sonication in 10 volumes of assay buffer (100 mM KCl/10 mM KH₂PO₄/K₂HPO₄, pH 7.5, at 4°C) with a protease inhibitor mixture (Complete Mini) and subjected to three centrifugation–resuspension cycles before being used for ligand binding assays.

[³H]Ro15-4513 Ligand Binding Assay. Membranes (or IP receptors bound to protein G–agarose beads) were resuspended in assay buffer (50 mM Tris-HCl, pH 8.0/1 M KCl/1 mM EDTA/0.5 mM DTT/2 mM benzamide/0.01% bacitracin/0.3 mM PMSF/10 $\mu\text{g}/\text{ml}$ trypsin inhibitor) by sonication. Resuspended membranes were incubated (in a volume of 0.5 ml) for 60 min on ice in the presence of [³H]Ro15-4513 (33.3 Ci/mmol, PerkinElmer) and various concentrations of competing ligands. Membranes (10–40 μg of protein per filter) were collected by rapid filtration on GF/B filters (Whatman). After three washing steps with 10 ml of assay buffer, the filter-retained radioactivity was counted in a LS3800 liquid scintillation counter (Beckman). Nonspecific binding was determined in the presence of 10 μM Ro15-4513, and DZ-IS binding was determined in the presence of 10 μM DZ. Data for binding curves were fitted by using a nonlinear least-squares method with the following equations: $B(c) = B_{\text{max}}/(1 + (K_d/c)^n)$, for binding curves, and $B(c) = B_{\text{max}}/(1 + (c/IC_{50})^n)$, where c is the concentration of ligand, B is binding, B_{max} is maximal binding, K_d is the dissociation constant, n is the Hill

